

Brand name: Zenol Xum SX

CHARACTERISTICS

Second-class OTC drug

●Topical analgesic anti-inflammatory agent

Felbinac penetrates into the affected area, and helps relieve joint, muscle, and shoulder pains.

This is a stick which is easily applied to the affected area without staining hands.

As it is not sticky, it allows putting on clothes immediately after use.

Easy-to-use new designed container



ACTIVE INGREDIENTS

Per 100 g

ACTIVE INGREDIENTS	Content
Felbinac	3.0 g
L-Menthol	3.0 g

Inactive ingredients: Isopropanol, Propylene Glycol, Macrogol, Polyoxyethylene Hardened Caster Oil, Sodium Stearate, 1,3-Butylene Glycol, Oleyl Alcohol, Squalane, pH Adjuster

For further information:

Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
Manufactured and Distributed by:	Mikasa Seiyaku Co., Ltd.
Dosage form	Stick type agent
Packaging unit	43 g x 1 box
Manufacturer's recommended price	1,500 yen
JAN Code	4987117372014
Expiration	3 years

INDICATIONS

Joint pain, muscle pain, shoulder pain due to stiff shoulder, low back pain, bruise, sprain, tenosynovitis (pain and swelling in hands, wrists, and ankles), elbow pain (tennis elbow, etc.)

DOSAGE AND ADMINISTRATION

Apply an appropriate amount to the affected area twice to four times daily.

<Precautions>

- (1) Strictly follow the dosing instructions.
 - (2) Be careful not to get the agent into the eyes. If the agent gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist.
 - (3) For external use only. Do not take the agent orally.
 - (4) Do not cover the affected area with a poorly-ventilated material such as wrap film, etc. after applying the agent.
 - (5) Wipe off sweat on the affected area before use.
 - (6) Twist the container until 4 to 5 mm of agent appears before use, to prevent the container from directly touching the affected area.
 - (7) Stop using the agent when the bottom of the container appears and the upper side of the bottom is exposed.
- Do not forcibly use the agent remaining in the bottom of the container.

PRECAUTIONS FOR USE

The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

1. The following persons should not use the agent.

- (1) Persons with a history of allergic symptoms to the agent or any of the ingredients of the agent
- (2) Persons with a history of asthma
- (3) Pregnant women or possibly pregnant women
- (4) Children under 15 years

2. Do not apply to the following areas.

- (1) Around the eyes, mucous membranes, etc.
- (2) Eczema, contact dermatitis, wounds
- (3) Athlete's foot (tinea pedis), tinea cruris, etc. or purulent areas

 Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before using the agent.

- (1) Persons under treatment by a physician
- (2) Persons with a history of allergic symptoms to any drug

2. The following symptoms may be adverse reactions to the agent. If any of these symptoms occur, discontinue the use of the agent immediately, and consult a physician, pharmacist or registered distributor with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching, swelling, tingling sensation, contact dermatitis

The following serious symptom may rarely occur. If the symptom occurs, consult a physician immediately.

Symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, larynx itching, difficulty breathing, palpitation, consciousness clouding, etc. occur immediately after using the agent.

3. If symptoms do not improve after using the agent for about one week, discontinue the use of the agent, and consult a physician, pharmacist or registered distributor with this leaflet.