Brand name: Zenol Tick E

RISK CATEGORIES

Third-class OTC drug

Topical analgesic anti-inflammatory agent

CHARACTERISTICS

 Methyl salicylate is effective for stiff shoulder and muscle fatigue by penetrating into the affected areas.

This is a stick which is easily applied to the affected area without staining hands. As it is not sticky, it allows putting on clothes immediately after use.

It has a cool flavor.





ACTIVE INGREDIENTS

• Per 100 g

For further information:

		Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
ACTIVE INGREDIENTS	Content	Manufactured and Distributed by:	Mikasa Seiyaku Co., Ltd.
Methyl Salicylate	17.50 g	Dosage form	Stick type agent
L-Menthol	6.00 g	Packaging unit	33 g x 1 box
dl-Camphor	5.00 g	Manufacturer's suggested retail price	JPY 935 (JPY 850 excluding tax)
Glycyrrhetinic Acid	0.02 g	JAN Code	4987117372052
Inactive ingredients: Isopropanol Flavor, Sodium Stearate, Glycerin, Caster Oil		Expiration	3 years

DOSAGE AND ADMINISTRATION

Usually, apply an appropriate amount to the affected area once to several times daily.

INDICATIONS

Stiff shoulder, muscle fatigue, muscle pain, low back pain, joint pain, bruise, sprain, frostbite

<Pre><Pre>cautions>

- (1) Strictly follow the dosing instructions.
- (2) Be careful not to get the agent into the eyes. If the agent gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist.
- (3) For children, use under the supervision of their guardians.
- (4) For external use only. Do not take the agent orally.
- (5) Wipe off sweat on the affected area before use.
- (6) Twist the container until 4 to 5 mm of agent appears before use, to prevent the container from directly touching the affected area.
- (7) Stop using the agent if the bottom of the container appears and becomes unstable.

Do not forcibly use the agent remaining in the bottom of the container.

PRECAUTIONS FOR USE

▼The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.) Do not apply to the following areas.

(1) Around the eyes, mucous membranes, etc.

(2) Eczema, contact dermatitis, wounds

Consultation

 The following persons should consult a physician, pharmacist or registered salesperson before using the agent.

Persons with a history of allergic symptoms due to any drug

The following symptoms may be adverse reactions to the agent. If any of these symptoms occur, discontinue
the use of the agent immediately, and consult a physician, pharmacist or registered salesperson with this
leaflet.

Affected body system	Symptoms	
Skin	Rash, redness, itching	

3. If symptoms do not improve after using the agent for five to six days, discontinue the use of the agent, and consult a physician, pharmacist or registered salesperson with this leaflet.

Precautions in storage and handling

- (1) Do not bring the product close to fire. And do not throw the used product into fire.
- (2) Store in a cool place, away from direct sunlight.
- (3) Keep out of the reach of children.
- (4) As the product is volatile, screw the cap tightly after use and avoid exposure to the outer air.
- (5) Do not transfer to other container. (It may cause misuse or change the quality.)
- (6) Take care as the product adhered to the painted matters (furniture, floor, accessory, etc.), artificial fiber, silk, leather, etc. may change their properties.
- (7) Do not use the product past the expiration date. After opening, use as soon as possible even before the expiration date to retain the quality.

Contact Information

If you have any questions regarding drug, please contact the shop where you purchased the product or the shop below.

Distributor

Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

₹101-8444 1-27 Kanda Nishiki-cho, Chiyoda-ku, Tokyo

Phone: 120-4527-66

Acceptance time 9:00 to 17:00 (excluding Saturdays, Sundays, and holidays)

Website https://www.taiho.co.jp/

Manufactured and marketed by Mikasa Pharmaceutical Co.,Ltd. 2-3-1 Toyotama Kita, Nerima-ku, Tokyo