

Brand name: Tiovita Drink 3000

CHARACTERISTICS
 Designated quasi-drug
 ● Vitamin health drink
 - The amount of taurine contained is the greatest ever since Tiovita Drink was released.
 - The medicine contains 3,000 mg of taurine, vitamin B groups, and carnitine hydrochloride as active ingredients.
 - Easy-to-take "mango flavor"



ACTIVE INGREDIENTS
 Per 100 mL bottle
 Taurine 3,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B1) 10 mg, Vitamin B2 Sodium Phosphate 10 mg, Vitamin B6 10 mg, Carnitine Hydrochloride 100 mg, Anhydrous Caffeine 50 mg
 Inactive ingredients: White Soft Sugar, Glucose, DL- Malic Acid, Propyl Gallate, Sodium Benzoate, Paraben, pH Adjuster, Vanillin, Flavor
 ● Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about.

For further information:

| | | | | |
|----------------------------------|--------------------------------|--------------------|---------------------|---------------------|
| Manufactured and Distributed by: | TAIHO PHARMACEUTICAL CO., LTD. | | | |
| Dosage form | Liquid | | | |
| Packaging unit | 100 mL x 1 bottle | 100 mL x 3 bottles | 100 mL x 10 bottles | 100 mL x 50 bottles |
| Manufacturer's recommended price | 220 yen | 660 yen | 2,200 yen | 11,000 yen |
| JAN Code | 4987117438512 | 4987117438574 | 4987117438543 | 4987117438604 |
| Expiration | 3 years | | | |

DOSAGE AND ADMINISTRATION
 Adults (15 years and over): Take 1 bottle (100 mL) once daily. ● Do not use in children (under 15 years). ● Follow the dosing instruction. (Please pay attention to overdose if other products containing vitamins are concomitantly used.)

INDICATIONS
 ○ Nutritional fortification ○ Weak constitution ○ Supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation

PRECAUTIONS FOR USE
 ■ Consultation
 1. In the following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine. (1) If rash occurs after using the medicine. (2) If symptoms do not improve after using the medicine for a while.
 2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.